

**INTERNATIONAL AS
MATHEMATICS**

9665

FM02 Further Pure Mathematics Unit 2

Mark scheme

January 2019

Version: 1.0 Final

Mark schemes are prepared by the Lead Assessment Writer and considered, together with the relevant questions, by a panel of subject teachers. This mark scheme includes any amendments made at the standardisation events which all associates participate in and is the scheme which was used by them in this examination. The standardisation process ensures that the mark scheme covers the students' responses to questions and that every associate understands and applies it in the same correct way. As preparation for standardisation each associate analyses a number of students' scripts. Alternative answers not already covered by the mark scheme are discussed and legislated for. If, after the standardisation process, associates encounter unusual answers which have not been raised they are required to refer these to the Lead Assessment Writer.

It must be stressed that a mark scheme is a working document, in many cases further developed and expanded on the basis of students' reactions to a particular paper. Assumptions about future mark schemes on the basis of one year's document should be avoided; whilst the guiding principles of assessment remain constant, details will change, depending on the content of a particular examination paper.

Further copies of this mark scheme are available from oxfordaqaexams.org.uk

Q	Answer	Mark	Comments
1	$hf(x, y) = \frac{0.1}{1.2^2 + 0.8^2}$ $= 0.048077$	M1 A1	PI
	$y_2 = 0.8 + 0.048077 = 0.848077$	m1	0.8+ their value of $hf(x, y)$
	$y_3 = 0.848077 + \frac{0.1}{1.3^2 + 0.848077^2}$ (= 0.88958)	m1	
	0.890	A1	CAO – candidate's final answer
	Total	5	

Q	Answer	Mark	Comments
2	[f(3) =] -9 and [f(4) =] 20	B1	PI
	$x_1 = 3 + \frac{9}{20 + 9}$	M1	PI
	$x_1 = \frac{96}{29}$	A1	
	$f(x_1) [= -2.2067] < 0$	B1	
	$f(x_1) < 0$ so $\alpha > \frac{96}{29}$ and $\frac{96}{29} < \alpha < 4$	E1	
	Total	5	


	Answer	Mark	Comments
3(a)	$y = ax^n \Rightarrow \log_{10} y = \log_{10} ax^n$ $\log_{10} y = \log_{10} a + \log_{10} x^n$	M1	Take logs and apply one log law correctly PI
	$\log_{10} y = \log_{10} a + n\log_{10} x$	m1	Apply a further log law correctly.
	$Y = \log_{10} a + nX$ (which is a linear relationship between Y and X.)	A1	Correct eqn. with base 10 (or lg or later evidence of use of base 10 if log without base here)
3(b)	0.602, 0.778, 0.903	B1	At least 2 sig fig
	1.96, 2.37, 2.66	B1	At least 2 sig fig
3(c)	Their four points plotted correctly	B1ft	
	Straight line drawn through their points	B1ft	
3(d)	$a = 10^{0.58}$	M1	10 to the power of their intercept
	$a = 3.8$	A1	[3.3, 4.2]
	Gradient found	M1	
	$n = 2.3$	A1	[2.1, 2.5]
	Total	11	

Q	Answer	Mark	Comments
4(a)	$\begin{bmatrix} 3 & 0 \\ 0 & 1 \end{bmatrix}$	B1	
4(b)	W in correct place $(-3, -3), (-1, -3)$ and $(-3, -6)$	B1 B1	Two vertices correct B1B0
4(c)	Method 1		
	$\det(\mathbf{P}) = 3$	B1ft	
	$\det(\mathbf{Q}) = -1$	B1	PI
	$\det(\mathbf{QP}) = -3$	B1	
	Alternative method		
	$\mathbf{Q} = \begin{bmatrix} 0 & -1 \\ -1 & 0 \end{bmatrix}$	B1	
	$\mathbf{QP} = \begin{bmatrix} 0 & -1 \\ -3 & 0 \end{bmatrix}$	M1	
	$\det(\mathbf{QP}) = -3$	A1	
	Total	6	

Q	Answer	Mark	Comments
5(a)	Method 1		
	$\det(\mathbf{B}) = 2$	M1	
	$\mathbf{B}^{-1} = \begin{bmatrix} 1 & 1 \\ -\frac{1}{2} & -\frac{1}{2} \\ \frac{1}{2} & -\frac{1}{2} \end{bmatrix}$	A1	
	$\mathbf{A} = \mathbf{CB}^{-1}$	M1	
	$\mathbf{A} = \begin{bmatrix} 1 & -3 \\ 2 & -4 \end{bmatrix}$	A1	
	Alternative method		
	Let $\mathbf{A} = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$ then $\begin{bmatrix} a & b \\ c & d \end{bmatrix} \begin{bmatrix} -1 & 1 \\ -1 & -1 \end{bmatrix} = \begin{bmatrix} 2 & 4 \\ 2 & 6 \end{bmatrix}$	M1	
	$\begin{aligned} -a - b &= 2 \\ a - b &= 4 \\ -c - d &= 2 \\ c - d &= 6 \end{aligned}$	M1	For four equations
	$a = 1$ and $b = -3$ or $c = 2$ and $d = -4$	M1	For solving one pair of equations
	$\mathbf{A} = \begin{bmatrix} 1 & -3 \\ 2 & -4 \end{bmatrix}$	A1	

5(b)	$\mathbf{B}^2 = \begin{bmatrix} 0 & -2 \\ 2 & 0 \end{bmatrix}$	M1	
	$\mathbf{B}^4 = (\mathbf{B}^2)^2 = \begin{bmatrix} -4 & 0 \\ 0 & -4 \end{bmatrix}$	M1 A1	
	$\mathbf{B}^4 = k\mathbf{I}$ where $k = -4$	E1	
5(c)(i)	Scale factor = $\sqrt{2}$	B1	
5(c)(ii)	-135°	B1	Or 135° clockwise oe. Do not accept just 135°
5(d)	$\mathbf{B}^{21} = (\mathbf{B}^4)^5 \times \mathbf{B}$	M1	
	$= (-4)^5 \times \mathbf{B}$ or $-1024\mathbf{B}$	M1	
	$\mathbf{B}^{21} = \begin{bmatrix} 1024 & -1024 \\ 1024 & 1024 \end{bmatrix}$	A1	
	Total	13	

Q	Answer	Mark	Comments
6(a)	$\frac{2}{3}$	B1	
6(b)	$G_X(t) = \sum_{n=1}^3 t^n \left(\frac{1}{3}\right)$	M1	Applies formula for $G_X(t)$
	$= \frac{1}{3}(t + t^2 + t^3)$ oe	A1	Ignore subsequent incorrect working
6(c)	Alternative Method 1		
	$G_{X+Y}(t) = \frac{1}{3}(t + t^2 + t^3)(0.6 + 0.4t)$	M1	Multiplies their $G_X(t)$ and $G_Y(t)$
	$\frac{1}{3}t \times 0.6$	M1	Multiplies out the required terms to find the coefficient of t Implied by correct answer
	$= 0.2$	A1	Accept 1/5 oe
	Alternative method 2		
	Y Bernoulli, $p = 0.4$ or $Y \sim B(1, 0.4)$	M1	Identifies distribution of Y
	$P(X + Y = 1)$ means $X = 1$ and $Y = 0$ or $P(X + Y = 1) = P(X = 1)P(Y = 0)$	M1	Identifies possible combinations of X and Y
	$= \frac{1}{3}(1 - 0.4) = 0.2$	A1	Accept 1/5 oe
	Total	6	


Q	Answer	Mark	Comments
7(a)	$(1 - 0.6)^3 = 0.064$	B1	Accept 8/125 oe
7(b)	$P(W H) = \frac{P(H W)P(W)}{P(H)}$ $= \frac{0.01 \times 0.7}{0.064}$	M1	Applies Bayes Theorem to find P(W H)
	$= \frac{7}{64} \text{ or AWRT } 0.109$	A1ft	ft their P(H) from (a) provided $0 < P(H) < 1$
	$P(H' \cap W) = P(W) - P(H \cap W)$ $P(H')P(W H') = P(W) - P(H)P(W H)$ $(1 - 0.064)P(W H') = 0.7 - 0.064 \times \frac{7}{64}$	M1	Method to find P(W H')
	$P(W H') = \frac{77}{104} \text{ or } 0.740 \text{ AWRT}$	A1ft	ft their P(H) from (a) provided $0 < P(H) < 1$
		A1ft	ft their probabilities Need to score at least one M1
	Total	6	

Q	Answer	Mark	Comments
8(a)	$E(2P^2 - 5) = 2E(P^2) - 5$ or $E(PQ) = 2E(P^3) - 5E(P)$	M1	One correct formula seen
	$E(2P^2 - 5) = 2 \times 5 - 5 = 5$ or $E(PQ) = 2 \times 14.6 - 5 \times 2 = 19.2$	A1	One correct value Accept 96/5 oe for 19.2 Can be implied by correct final answer
	$E(2P^2 - 5) = 2 \times 5 - 5 = 5$ and $E(PQ) = 2 \times 14.6 - 5 \times 2 = 19.2$	A1	Both correct values Accept 96/5 oe for 19.2 Can be implied by correct final answer
	$\text{Cov}(P, Q) = E(PQ) - E(P)E(Q)$ $= 19.2 - 2 \times 5$	M1	Applies formula for Cov (P, Q)
	$= 9.2$	A1	Accept 46/5 oe
8(b)	$\text{Var}(P) = E(P^2) - (E(P))^2$ $= 5 - 2^2$ $= 1$	B1	Finds Var (P) Can be implied by correct final answer
	$\text{Var}(P + Q)$ $= \text{Var}(P) + \text{Var}(Q) + 2 \text{Cov}(P, Q)$ $= 1 + 8 + 2 \times 9.2$	M1	Applies formula for Var (P + Q)
	$= 27.4$	A1	Accept 137/5 oe
	Total	8	

Q	Answer	Mark	Comments
9	$MLT^{-2} = [k]L^{\frac{3}{2}}T^{-\frac{3}{2}}$ $[k] = ML^{-\frac{1}{2}}T^{-\frac{1}{2}}$	M1 A1	
	Total	2	

Q	Answer	Mark	Comments
10(a)	$I = \frac{1}{2} \times 2500 \times 10$ $= 12500 \text{ Ns}$	B1	
10(b)	$-12500 = 2000U - 2000 \times 20$ $U = \frac{2000 \times 20 - 12500}{2000}$ $U = 13.75$	M1 A1 A1	M1: Equation with correct terms but any signs. A1: Correct equation. A1: Correct U .
	Total	4	

Q	Answer	Mark	Comments
11(a)	$5mU + 4mU = mv_A + 4mv_B$ $9U = v_A + 4v_B$ $v_A - v_B = -e(5U - U)$ $v_A - v_B = -4eU$ $9U = v_B - 4eU + 4v_B$ $v_B = \frac{U(9 + 4e)}{5}$	M1 A1 M1 A1 A1	
11(b)	$v_A = \frac{U(9 + 4e)}{5} - 4eU$ $v_A = \frac{U(9 - 16e)}{5}$ $9 - 16e < 0$ $e > \frac{9}{16}$	B1 M1 A1	
	Total	8	

Q	Answer	Mark	Comments
12	$v_{MY}^2 = 8^2 + 5^2 - 2 \times 8 \times 5 \cos 30^\circ$ $v_{MY}^2 = 89 - 40\sqrt{3}$ $\frac{\sin \theta}{5} = \frac{\sin 30^\circ}{\sqrt{89 - 40\sqrt{3}}}$ $\theta = \sin^{-1} \frac{5}{2\sqrt{89 - 40\sqrt{3}}}$ $\theta = 34.26357 \dots^\circ$ <p>Min Sep = $2000 \times \sin(60 - \theta) = 868 \text{ m}$</p>	<p>M1</p> <p>A1</p> <p>M1</p> <p>A1</p> <p>M1</p> <p>A1</p>	<p>Accept 869</p>
	Additional Guidance		
	 <p>The diagram shows a vector triangle. A vertical vector of length 8 m s^{-1} is shown. A dashed line is drawn from the tip of this vector to a point on a horizontal line of length 2000 m. The angle between the vertical vector and the dashed line is θ. A second vertical vector of length 5 m s^{-1} is shown perpendicular to the dashed line, starting from the tip of the 8 m s^{-1} vector.</p>		
Total		6	