

Please write clearly in block capitals.

Centre number 

--	--	--	--	--

Candidate number 

--	--	--	--

Surname \_\_\_\_\_

Forename(s) \_\_\_\_\_

Candidate signature \_\_\_\_\_

I declare this is my own work.

# INTERNATIONAL A-LEVEL MATHEMATICS

(9660/MA03) Unit P2 Pure Mathematics

Thursday 14 January 2021 07:00 GMT Time allowed: 2 hours 30 minutes

## Materials

- For this paper you must have the Oxford International AQA Booklet of Formulae and Statistical Tables (enclosed).
- You may use a graphical calculator.

## Instructions

- Use black ink or black ball-point pen. Pencil should only be used for drawing.
- Fill in the boxes at the top of this page.
- Answer **all** questions.
- You must answer the questions in the spaces provided. Do not write outside the box around each page or on blank pages.
- If you need extra space for your answer(s), use the lined pages at the end of this book. Write the question number against your answer(s).
- Do all rough work in this book. Cross through any work you do not want to be marked.

## Information

- The marks for questions are shown in brackets.
- The maximum mark for this paper is 120

## Advice

- Unless stated otherwise, you may quote formulae, without proof, from the booklet.
- Show all necessary working; otherwise marks may be lost.

For Examiner's Use	
Question	Mark
1	
2	
3	
4	
5	
6	
7	
8	
9	
10	
11	
12	
13	
<b>TOTAL</b>	



J A N 2 1 M A 0 3 0 1

Answer **all** questions in the spaces provided.

- 1** The functions  $f$  and  $g$  are defined with their respective domains by

$$f(x) = x - 5 \quad \text{for all real values of } x$$

$$g(x) = \frac{25}{x+4} \quad \text{for all real values of } x, \quad x \neq -4$$

The composite function  $fg$  is denoted by  $h$

- 1 (a)** Find  $h(x)$  giving your answer as a single fraction.

**[2 marks]**

---

---

---

---

---

---

---

Answer \_\_\_\_\_



**1 (b)** The inverse of  $h$  is  $h^{-1}$

**1 (b) (i)** Find  $h^{-1}(x)$

**[3 marks]**

---

---

---

---

---

---

---

---

---

---

---

---

Answer \_\_\_\_\_

**1 (b) (ii)** Find the range of  $h^{-1}$

**[1 mark]**

---

---

---

---

Answer \_\_\_\_\_



2 The line  $l_1$  has equation  $\mathbf{r} = \begin{bmatrix} 4 \\ -2 \\ -3 \end{bmatrix} + \lambda \begin{bmatrix} -1 \\ 5 \\ 2 \end{bmatrix}$

The line  $l_2$  has equation  $\mathbf{r} = \begin{bmatrix} -1 \\ 5 \\ 11 \end{bmatrix} + \mu \begin{bmatrix} -1 \\ -4 \\ c \end{bmatrix}$

2 (a) In the case where  $l_1$  and  $l_2$  intersect, find

2 (a) (i) the value of  $c$

[3 marks]

---

---

---

---

---

Answer \_\_\_\_\_

2 (a) (ii) the coordinates of the point of intersection.

[1 mark]

Answer \_\_\_\_\_

2 (b) In the case where  $l_1$  and  $l_2$  are perpendicular, find the value of  $c$

[3 marks]

---

---

---

---

---

Answer \_\_\_\_\_



**3** It is given that  $y = 3\sin\theta - 3\cos\theta$

**3 (a)** Express  $y$  in the form  $R\sin(\theta - \alpha)$  where  $R$  is a surd and  $0^\circ < \alpha < 90^\circ$

**[2 marks]**

---

---

---

---

Answer \_\_\_\_\_

**3 (b)** Hence find

**3 (b) (i)** the greatest value of  $y^2$

**[1 mark]**

---

Answer \_\_\_\_\_

**3 (b) (ii)** the least value of  $y^2$

**[1 mark]**

---

Answer \_\_\_\_\_

**3 (b) (iii)** the values of  $\theta$  in the interval  $-90^\circ < \theta < 90^\circ$  for which  $y = -\frac{3\sqrt{6}}{2}$

**[3 marks]**

---

---

---

---

---

---

---

---

---

---

Answer \_\_\_\_\_



- 4 (a)** Describe a sequence of **two** geometrical transformations that maps the graph of  $y = \cos x$  onto the graph of  $y = 1 + 2\cos x$

[4 marks]

---

---

---

---

---

---

---

---

---


---

- 4 (b)** Sketch the graph of the curve with equation

$$y = 1 + 2\cos x \quad \text{for} \quad -\pi < x < \pi$$

indicating the value of  $y$  where the curve crosses the  $y$ -axis.

[2 marks]



- [6 marks]**

[illegible]

Answer \_\_\_\_\_

12

**Turn over ►**



**[2 marks]**

---

---

---

---

---

$$\int_0^{0.5} \sqrt{(1+x^2)} \, dx$$

**[4 marks]**

This image shows a single sheet of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.

Answer





$$\int_0^{0.5} \sqrt{(1+x^2)} \, dx$$

**[4 marks]**

[illegible]

Answer

10

**Turn over ►**



**6** It is given that  $\alpha + \beta = 45^\circ$  where  $\alpha$  and  $\beta$  are both positive.

**6 (a)** Find  $\tan \beta$  in terms of  $\tan \alpha$

**[2 marks]**

---

---

---

---

---

---

---

---

---

---

Answer \_\_\_\_\_

**6 (b)** Show that  $(1 + \tan \alpha)(1 + \tan \beta) = 2$

**[2 marks]**

---

---

---

---

---

---

---

---



**6 (c)** Find the exact value of  $\tan 22.5^\circ$

**[3 marks]**

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

Answer \_\_\_\_\_

7

**Turn over for the next question**

**Turn over ►**



The curve intersects the line  $y = 3 - 4x$  at a single point where  $x = \alpha$

**[2 marks]**

---

---

---

$$x_{n+1} = \frac{3 - \sin(\ln(2x_n))}{4}$$

Give your answers to three decimal places.

**[2 marks]**

---

$$x_2 = \quad \quad \quad x_3 =$$

Find the coordinates of a stationary point of the curve.

**[3 marks]**

---

---

---

---

---

Answer



Show that  $x^2 \frac{d^2 y}{dx^2} + x \frac{dy}{dx} + y = 0$

[illegible]

8 (a) Given that  $\cot x = \frac{\cos x}{\sin x}$ , use the quotient rule to show that

$$\frac{d}{dx}(\cot x) = -\operatorname{cosec}^2 x$$

[2 marks]

---

---

---

---

---

---

---

---

---

---

---

---



**8 (b) (i)** Find  $\frac{dx}{dy}$  giving your answer in terms of  $y$

---

---

---

---

---

Answer

**[3 marks]**

[illegible]

Answer

7



9 (a) Given that  $x^3 + y^3 = 3xy$  show that  $\frac{dy}{dx} = \frac{y - x^2}{y^2 - x}$

[2 marks]

---

---

---

---

---

---

---

---

9 (b) A curve has the equation  $x^3 + y^3 = 3xy$

9 (b) (i) Find the coordinates of the stationary point of the curve in the interval  $0 < x < 2^{\frac{2}{3}}$

Give your answer in exact form.

[3 marks]

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

Answer \_\_\_\_\_





**[4 marks]**

[illegible]

Answer \_\_\_\_\_

9



When  $t = 5$  the temperature of the object is  $70^\circ\text{C}$

**[1 mark]**

**10 (b)** Find the temperature of the object when  $t = 15$  giving your answer to one decimal place  
[6 marks]



**10 (c)** Find the value of  $t$  when the temperature of the object is  $40^{\circ}\text{C}$  giving your answer to one decimal place.

Answer \_\_\_\_\_

9



**[5 marks]**

[illegible]

Answer

$$\int_8^{15} \frac{\sqrt{x+1}}{x-3} dx$$

**[9 marks]**

[illegible]

[illegible]

Answer \_\_\_\_\_

14

**[4 marks]**

This image shows a blank sheet of white paper with horizontal ruling lines. The lines are evenly spaced and extend across the width of the page. There are no margins, text, or other markings on the paper.

Answer

**[2 marks]**

---

---

---

---

---

Answer



**12 (c)**

for small values of  $x$ , stating the rational values of  $D$ ,  $E$  and  $F$

**[4 marks]**

[illegible]

10

**Turn over ►**



13

$$x = ct, \quad y = \frac{c}{t} \quad \text{where } t > 0 \text{ and } c \text{ is a constant.}$$

The tangent at the point  $P\left(cp, \frac{c}{p}\right)$  on the curve meets the  $x$ -axis at  $A$  and the  $y$ -axis at  $B$

The normal at the point  $P$  meets the line  $y = x$  at  $C$  and the line  $y = -x$  at  $D$

**13 (a)**

**[1 mark]**

Answer

**13 (b)**

**[7 marks]**





**[3 marks]**

11



**There are no questions printed on this page**

*Do not write  
outside the  
box*

**DO NOT WRITE ON THIS PAGE  
ANSWER IN THE SPACES PROVIDED**





2 1 1 X M A 0 3

