

INTERNATIONAL A-LEVEL FURTHER MATHEMATICS FM04

(9665/FM04) Unit FS2 Statistics

Mark scheme

January 2023

Version: 1.0 Final



Mark schemes are prepared by the Lead Assessment Writer and considered, together with the relevant questions, by a panel of subject teachers. This mark scheme includes any amendments made at the standardisation events which all associates participate in and is the scheme which was used by them in this examination. The standardisation process ensures that the mark scheme covers the students' responses to questions and that every associate understands and applies it in the same correct way. As preparation for standardisation each associate analyses a number of students' scripts. Alternative answers not already covered by the mark scheme are discussed and legislated for. If, after the standardisation process, associates encounter unusual answers which have not been raised they are required to refer these to the Lead Examiner.

It must be stressed that a mark scheme is a working document, in many cases further developed and expanded on the basis of students' reactions to a particular paper. Assumptions about future mark schemes on the basis of one year's document should be avoided; whilst the guiding principles of assessment remain constant, details will change, depending on the content of a particular examination paper.

Further copies of this mark scheme are available from oxfordaqaexams.org.uk

Copyright information

OxfordAQA retains the copyright on all its publications. However, registered schools/colleges for OxfordAQA are permitted to copy material from this booklet for their own internal use, with the following important exception: OxfordAQA cannot give permission to schools/colleges to photocopy any material that is acknowledged to a third party even for internal use within the centre.

Copyright © 2023 Oxford International AQA Examinations and its licensors. All rights reserved.

Key to mark scheme abbreviations

м	Mark is for method
m	Mark is dependent on one or more M marks and is for method
Α	Mark is dependent on M or m marks and is for accuracy
В	Mark is independent of M or m marks and is for method and accuracy
Е	Mark is for explanation
\checkmark or ft	Follow through from previous incorrect result
CAO	Correct answer only
CSO	Correct solution only
AWFW	Anything which falls within
AWRT	Anything which rounds to
ACF	Any correct form
AG	Answer given
SC	Special case
oe	Or equivalent
A2, 1	2 or 1 (or 0) accuracy marks
<i>–x</i> EE	Deduct <i>x</i> marks for each error
NMS	No method shown
PI	Possibly implied
SCA	Substantially correct approach
sf	Significant figure(s)
dp	Decimal place(s)

Q	Answer	Marks	Comments
1(a)	$\chi^2 = \frac{s^2}{\sigma_0^2} \times (n-1) = \frac{100}{\sigma_0^2} \times 9$	M1	Use of correct statistic. PI Allow <i>n</i> for $n - 1$
	$\chi_9^2(0.975) = 19.023$	B1	Finds critical value
	$\sigma_0^2 > \frac{900}{19.023}$ [= 47.3111]	M1	Allow either >, ≥ or = oe
	$\sigma_0 = 6.878[31]$	A1	Must show answer at least 4 sf or explicitly state as 6.88 to 3 sf AG
		4	

Q	Answer	Marks	Comments
1(b)	$\sigma_0^2 < \frac{900}{\chi_9^2(0.025)} = \frac{900}{2.700} [= 333.333]$	M1	Allow either <, ≤ or = oe ft their degrees of freedom in (a)
	σ_0 =18.3	A1	AWRT Allow truncation to 18.2
		2	
	Question 1 Total	6	

Q	Answer					Marks	Comments
2(a)	$3\times0.7^2\times0.3$ or 0.3^3					M1	PI or one value (of 0.441 or 0.027) correct
	v P(V=v)	15 0.343	60 0.441	105 0.189	150 0.027	B1 A1	Both 60 and 105 needed Both 0.441 and 0.027 needed
						3	

Q	Answer	Marks	Comments
2(b)(i)	0.343 + 0.441[= 0.784] or $0.189 + 0.027[= 0.216]$ m 5 50	M1	PI ft their 0.441 or 0.027
	P(M=m) 0.784 0.216	A1	
		2	

Q	Answer	Marks	Comments
2(b)(ii)	E(M)=0.784 × 5 + 0.216 × 50[=14.72] or E(M ²)=0.784 × 5 ² + 0.216 × 50 ² [=559.6]	М1	PI ft their (b)(i)
	$Var(M) = 559.6 - 14.72^2$ Var(M) = 343	M1 A1	Use of Var $(M) = E(M^2) - (E(M))^2$ PI AWRT
		3	

Question 2 Total	8	
------------------	---	--

Q	Answer	Marks	Comments
3(a)	$\frac{27.8+30.4}{2} = 29.1$	B1	
		1	

Q	Answer	Marks	Comments
3(b)	Critical value $z = (\pm)1.96(00)$	B1	AWRT 1.96
	$30.4 - 27.8 = 2.6 = 2 \times 1.96 \times \frac{\sqrt{6.6}}{\sqrt{n}}$	M1	Use of $\frac{\sqrt{6.6}}{\sqrt{n}}$ in an equation to find <i>n</i>
	$n = \frac{6.6 \times 1.96^2}{1.3^2} = 15.0027 \text{ so } 15$	A1	AG CSO Either value for <i>n</i> given to at least three significant figures or calculation for <i>n</i> with correct substitution must be seen
		3	

Q	Answer	Marks	Comments
3(c)	30 is in the confidence interval	B1	Condone use of "it" for 30
	Evidence that the target (of mean conference call of 30 minutes) has been met	E1	Must be in context
		2	

Q	Answer	Marks	Comments
3(d)	It is a normal distribution with known [population] variance	B2	1 mark for each feature (normal distribution, known variance)
		2	

Question 3 To	I 8	
---------------	-----	--

Q	Answer	Marks	Comments
4(a)	$M'_{Z}(t) = t e^{\frac{1}{2}t^{2}}$ $M'_{Z}(0) = 0 \times e^{0} = 0$ $M''_{Z}(t) = (1+t^{2}) e^{\frac{1}{2}t^{2}}$ $\sigma^{2} = M''_{Z}(0) - \mu^{2}$ $= 1-0 = 1$	M1	Allow $ate^{\frac{1}{2}t^2}$
	$M_{Z}'(0) = 0 \times e^{0} = 0$	Α1	
	$M_{Z}''(t) = (1+t^{2})e^{\frac{1}{2}t^{2}}$	М1	Of form $\left(a+bt^2\right)e^{\frac{1}{2}t^2}$ oe
	$\sigma^2 = M_Z'(0) - \mu^2$	M1	
	= 1-0 = 1	A1	
		5	

Q	Answer	Marks	Comments
4(b)	$M_X(t) = \mathrm{e}^{at} \times \mathrm{e}^{\frac{1}{2}(bt)^2}$	M1	Use of $M_X(t) = e^{at} \times M_Z(bt)$
	$M_X(t) = e^{at + \frac{1}{2}b^2t^2}$	A1	
		2	

Q	Answer	Marks	Comments
4(c)	$E(X) = a$ and $Var(X) = b^2$	B1	Both E (X) and Var (X) required
		1	

Q	Answer	Marks	Comments
4(d)	$e^{\mu t +}$ or $e^{+\frac{1}{2}\sigma^2 t^2}$	M1	
	$e^{\mu t + \frac{1}{2}\sigma^2 t^2}$	A1	
		2	

Question 4 Tot	10	
----------------	----	--

Q	Answer	Marks	Comments
5(a)	$E(\overline{X}) = \frac{n\lambda}{n} = \lambda \text{ and } E(\overline{Y}) = \frac{n \times 2\lambda}{n} = 2\lambda$	B1	Both. PI
	$E(S) = \frac{\lambda + 2\lambda}{3} = \lambda \text{ or } E(T) = 2\lambda - \lambda = \lambda$	M1	Either found
	$\mathrm{E}(S) = \lambda$ and $\mathrm{E}(T) = \lambda$ so estimators are unbiased	A1	Statement and both estimators correct
		3	

Q	Answer	Marks	Comments
5(b)	$\operatorname{Var}(S) = \left(\frac{1}{3}\right)^{2} \operatorname{Var}(\overline{X}) + \left(\frac{1}{3}\right)^{2} \operatorname{Var}(\overline{Y})$	M1	Correct expression for $Var(S)$ or $Var(T)$ May be seen in (c)
	$\operatorname{Var}(T) = \operatorname{Var}(\overline{Y}) + \operatorname{Var}(\overline{X})$		
	$\operatorname{Var}(S) = \frac{1}{9} \times \frac{\lambda}{n} + \frac{1}{9} \times \frac{2\lambda}{n} = \frac{\lambda}{3n}$ $\operatorname{Var}(T) = \frac{\lambda}{n} + \frac{2\lambda}{n} = \frac{3\lambda}{n}$	A1	PI May be seen in (c)
	$\operatorname{Var}(T) = \frac{\lambda}{n} + \frac{2\lambda}{n} = \frac{3\lambda}{n}$	A1	PI May be seen in (c)
	Relative Efficiency = $\frac{\frac{1}{\operatorname{Var}(S)}}{\frac{1}{\operatorname{Var}(T)}} = \frac{\frac{3n}{\lambda}}{\frac{n}{3\lambda}}$	M1	ft their Var(<i>S</i>) and Var(<i>T</i>) oe
	[Relative Efficiency] = 9 [which is not a function of n , so the efficiency is independent of n]	A1	Answer of 9 is sufficient for award of mark
		5	

Q	Answer	Marks	Comments
5(c)	$\operatorname{Var}(S) \to 0 \text{ or } \operatorname{Var}(T) \to 0 \text{ as } n \to \infty$	M1	Either may be shown from a function of n that tends to zero
	so estimators are consistent	A1	Conclusion required CSO
		2	

Question 5 T	tal 10	
--------------	--------	--

Q	Answer	Marks	Comments
6(a)	$\int_{100}^{t} -\frac{\pi}{200} \sin\left(\frac{\pi x}{100}\right) dx$	M1	Must have correct limits
	$= \left[\frac{1}{2}\cos\left(\frac{\pi x}{100}\right)\right]_{100}^{t}$	M1	Integrand of form $a\cos\left(\frac{\pi x}{100}\right)$ oe
	$=\frac{1}{2}\cos\left(\frac{\pi t}{100}\right) - \frac{1}{2}\cos\left(\frac{100\pi}{100}\right)$		
	$F(t) = \begin{cases} 0 & t < 100\\ \frac{1}{2}\cos\left(\frac{\pi t}{100}\right) + \frac{1}{2} & 100 \le t \le 200\\ 1 & t > 200 \end{cases}$	A1	AG must see intermediate line with values substituted into integrand Limits for <i>t</i> need to be shown
		3	

Q			Ans	wer			Marks	Comments
6(b)(i)	F(160) – F(140), F(180) – F(160) or F(200) – F(180) seen						M1	Ы
	Interval 100- 120- 140- 160- 180- 120 140 160 180 200							
	Sprints 164 430 532 430 164					164		
		Either of 430 or 164 seen All 532, 430, 164						Allow +/– 1 for both A marks
							3	

Q	Answer	Marks	Comments
6(b)(ii)	H_0 : Reaction times have the same distribution as T H_1 : Reaction times do not have the same distribution as T	B1	oe , eg H ₀ : Suggested model is appropriate, Athletics trainer's claim is valid (condone true), Data fits given distribution Both hypotheses
	$\sum \frac{(O-E)^2}{E} = \frac{(145-164)^2}{164} + \frac{(390-430)^2}{430} + \frac{(561-532'')^2}{532''} + \frac{(470-430'')^2}{430''} + \frac{(154-164'')^2}{164''}$	М1	
	= 11.8 A1ft		ft their (b)(i) given to 1 decimal place
	v=5-1= 4	B1	
	$\chi^2(0.99) = 13.277$	B1	
	11.8 < 13.277, Do not reject H₀	A1ft	
	Sufficient evidence to support the athletics trainer's claim	E1ft	Must not be definite; consistent with conclusion on ${\rm H_0}$
		7	

		13	Question 6 Total
--	--	----	------------------

Q	Answer	Marks	Comments
7(a)(i)	The test is a two-tailed test.	B1	
		1	

Q	Answer	Marks	Comments
7(a)(ii)	$z = \frac{53.4 - 45 - 10}{\sqrt{\left(\frac{6^2}{60} + \frac{4^2}{80}\right)}}$	M1 M1	Correct numerator Correct denominator
	= -1.788(85)	A1	AWRT –1.79 <i>p</i> = 0.0736
	$z_{\rm crit} = +/-1.9600$	B1	AWRT 1.96
	–1.7889 > –1.9600 Do not reject H ₀	A1ft	Follow through their z and $z_{\rm crit}$
	Sufficient evidence to suggest that the mean length of Galapagos penguins is 10 cm more than that of Fairy penguins	E1	Gives a conclusion in context based on a comparison of the correct test statistic and correct critical value Condone definite conclusion
		6	

Q	Answer	Marks	Comments
7(b)	The result is valid as the sample is sufficiently large to use a normal approximation for the mean (Central Limit Theorem)	E1	oe must clearly state validity with reason Condone "can use" oe
		1	

Question 7 1

Q	Answer	Marks	Comments
8	<i>z</i> =1.6449	B1	AWRT 1.645
	$\overline{X_{c}} = 100 + 1.6449 \times \frac{10}{\sqrt{30}}$ $\begin{bmatrix} = 103.00316 \implies \text{Acceptance region: } \overline{X} < 103 \end{bmatrix}$	M1	
	$P\left(\overline{X} < 103 \mu\right) \le 0.05$	m1	PI Condone < or = [μ is the population mean.]
	$103 < \mu - 1.6449 imes rac{10}{\sqrt{30}}$	m1	Condone = Dependent on all previous method marks
	$\mu > 106.0(031)$	A1	AG Strict inequality sign required
		5	

Question 8 Tot	I 5	
----------------	-----	--

Q			Ansv	wer			Marks	Comments
9(a)		1						
	Computer	1	2	3	4	5		
	Difference	-2.2	+8.1	a–113.5	-6.6	-2.5	M1	Attempt differences; allow 1 mistake,
	and							allow negative of table values PI
	Computer	6	7	8	9	10		
	Difference	-7.6	+0.1	+4.0	-4.2	+1.2		
	$\overline{d} = \frac{-123.}{10}$	$\frac{a^2+a}{0} =$	0.1 <i>a</i> –	-12.32			B1	Allow negative, 12.32 – 0.1 <i>a</i>
	$\sum d^2 = 213$ $\left(=a^2 - 227\right)$	7 <i>a</i> +130	095.36	5)			М1	Allow $a^2 - ba + c$, with b and c positive values
	$s^{2} = \frac{1}{10 - 1} (1)$ $= \frac{1}{9} (11577)$. ²)			
	$= 0.1a^2 - 22$	2.484 <i>a</i>	+1286	5.392 8			A1	ое
	$t = \frac{\overline{d}}{\left(\frac{s}{\sqrt{10}}\right)} =$	$=\frac{1}{\sqrt{0.1a}}$	$\frac{0}{a^2-22}$	$\frac{1a-12.3}{2.48\dot{4}a+1}$	2 286.3	<u>928</u>	M1	ft with their mean and variance Allow $-t$
	$t = \frac{\sqrt{10}}{\sqrt{0.1a^2}}$	(0.1 <i>a - 1</i> - 22.48	12.32) a+12	86			A1	AG Must be convincingly shown
							6	

Q	Answer	Marks	Comments
9(b)	$ \begin{array}{l} H_{0} \colon \ \mu_{new} = \mu_{old} \\ H_{1} \colon \ \mu_{new} < \mu_{old} \end{array} $	B1	oe
	<i>t</i> = –1.23(1)	M1	Correct substitution of $a = 91.8$ into formula Condone 1.23
	<i>v</i> =9	B1	Ы
	Critical value $t_9 = 1.383$	B1	
	-1.23 > -1.383 , Do not reject H ₀	A1ft	Allow 1.23 < 1.383 ft their t and critical value
	Insufficient evidence to support the reduction in start-up times	E1	Gives a conclusion in context based on a comparison of the correct test statistic and correct critical value
			Condone definite conclusion
		6	

Question 9 To
